



**UNIVERSIDADE ESTADUAL DE CAMPINAS**  
**FACULDADE DE ODONTOLOGIA DE PIRACICABA**

**ELIS JANAÍNA LIRA DOS SANTOS**

**ANÁLISE ULTRAESTRUTURAL DOS CEMENTÓCITOS SUBMETIDOS À  
APOSIÇÃO EXPERIMENTAL DO CEMENTO DENTAL**

**ULTRASTRUCTURAL ANALYSIS OF CEMENTOCYTES UNDER  
EXPERIMENTALLY-INDUCED DENTAL CEMENTUM APPPOSITION**

Piracicaba

2017

**ELIS JANAÍNA LIRA DOS SANTOS**

**ANÁLISE ULTRAESTRUTURAL DOS CEMENTÓCITOS SUBMETIDOS À  
APOSIÇÃO EXPERIMENTAL DO CEMENTO DENTAL**

**ULTRASTRUCTURAL ANALYSIS OF CEMENTOCYTES UNDER  
EXPERIMENTALLY-INDUCED DENTAL CEMENTUM APPPOSITION**

Dissertação de Mestrado apresentada à Faculdade de Odontologia de Piracicaba da Universidade Estadual de Campinas como parte dos requisitos exigidos para obtenção do título de Mestre em Clínica Odontológica, na Área de Periodontia.

Dissertation presented to the Piracicaba Dental School of the University of Campinas in partial fulfillment of the requirements for the degree of Master in Clinical Dentistry, in Periodontics area.

Orientador: Prof. Dr. Francisco Humberto Nociti Júnior

Coorientadora: Profa. Dra. Cristiane Ribeiro Salmon

ESTE EXEMPLAR CORRESPONDE À VERSÃO FINAL DA DISSERTAÇÃO DE MESTRADO DEFENDIDA PELA ALUNA ELIS JANAÍNA LIRA DOS SANTOS, E ORIENTADA PELO PROF. DR. FRANCISCO HUMBERTO NOCITI JÚNIOR.

Piracicaba

2017

**Agência(s) de fomento e nº(s) de processo(s):** FAPESP, 2015/06372-2; CAPES  
**ORCID:** <http://orcid.org/http://orcid.org/ht>

Ficha catalográfica  
Universidade Estadual de Campinas  
Biblioteca da Faculdade de Odontologia de Piracicaba  
Marilene Girello - CRB 8/6159

Sa59a Santos, Elis Janaína Lira dos, 1988-  
Análise ultraestrutural dos cementócitos submetidos à aposição experimental do cimento dental / Elis Janaína Lira dos Santos. – Piracicaba, SP : [s.n.], 2017.

Orientador: Francisco Humberto Nociti Júnior.  
Coorientador: Cristiane Ribeiro Salmon.  
Dissertação (mestrado) – Universidade Estadual de Campinas, Faculdade de Odontologia de Piracicaba.

1. Cimento dentário. 2. Cementogênese. 3. Microscopia. I. Nociti Júnior, Francisco Humberto, 1967-. II. Salmon, Cristiane Ribeiro, 1975-. III. Universidade Estadual de Campinas. Faculdade de Odontologia de Piracicaba. IV. Título.

Informações para Biblioteca Digital

**Título em outro idioma:** Ultrastructural analysis of cementocytes under experimentally-induced dental cementum apposition

**Palavras-chave em inglês:**

Dental cementum

Cementogenesis

Microscopy

**Área de concentração:** Periodontia

**Titulação:** Mestra em Clínica Odontológica

**Banca examinadora:**

Francisco Humberto Nociti Júnior [Orientador]

Bruna Rabelo Amorim

Karina Gonzales Silvério Ruiz

**Data de defesa:** 15-02-2017

**Programa de Pós-Graduação:** Clínica Odontológica



**UNIVERSIDADE ESTADUAL DE CAMPINAS**

**Faculdade de Odontologia de Piracicaba**



A Comissão Julgadora dos trabalhos de Defesa de Dissertação de Mestrado, em sessão pública realizada em 15 de Fevereiro de 2017, considerou a candidata ELIS JANAÍNA LIRA DOS SANTOS aprovada.

PROF. DR. FRANCISCO HUMBERTO NOCITI JÚNIOR

PROF<sup>a</sup>. DR<sup>a</sup>. BRUNA RABELO AMORIM

PROF<sup>a</sup>. DR<sup>a</sup>. KARINA GONZALES SILVERIO RUIZ

A Ata da defesa com as respectivas assinaturas dos membros encontra-se no processo de vida acadêmica do aluno.

## DEDICATÓRIA

Dedico este trabalho a Deus, que conduziu minhas escolhas até aqui. Também dedico à minha mãe, Risalva, meu pai, Aldeci, e irmã, Erika, por tornarem possível a substancialização deste sonho.

## AGRADECIMENTOS

À Deus, por guiar os meus passos, ainda que nem existisse chão, me fazendo acreditar que basta colocar um pé a frente do outro e confiar. Por mostrar o quanto não temos controle sobre a vida, tudo tem um tempo para acontecer. Por me fazer forte nos momentos de fraqueza, e segurar o sorriso em meu rosto ainda que o mundo parecesse desmoronar. Por inundar meu ser de otimismo e fé diante de todas as adversidades e colocar pessoas incríveis no meu caminho, ainda que eu não mereça. Por ser o Deus do impossível.

Aos meus estimados pais, José Aldeci dos Santos e Risalva Gomes de Lira Santos, por terem escolhido ser os melhores pais possíveis, por me trazer a este mundo e ensinar a viver nele. Pelo suporte desde sempre, pelo respeito, por todas as vezes que usaram a palavra 'não', por serem meu exemplo e inspiração. Tudo o que fizer sempre será pensando em retribuir o que me foi dado com tanto amor, tanto para eles como para o mundo.

À minha irmã Erika Janine Lira dos Santos, pelo companheirismo, por virar as madrugadas comigo para estudar ou finalizar trabalhos, por ter feito minha inscrição na seleção do mestrado, por compartilhar os pais, a casa, a vida. Por ser uma parte de mim, minha melhor amiga.

À Universidade Estadual de Campinas (UNICAMP), na pessoa do magnífico Reitor Prof. Dr. José Tadeu Jorge e à Faculdade de Odontologia de Piracicaba (FOP), por meio do Diretor Prof. Dr. Guilherme Elias Pessanha Henriques. À Prof<sup>a</sup>. Dr<sup>a</sup>. Karina Gonzales Silvério Ruiz, coordenadora do Programa de Pós-Graduação em Clínica Odontológica e à Prof<sup>a</sup> Dr<sup>a</sup> Cinthia P. M. Tabchoury, coordenadora do Programa de Pós-Graduação da FOP/UNICAMP.

Ao Prof. Dr. Francisco Humberto Nociti Júnior, pesquisador notável, com sede de conhecimento e dotado de inteligência singular, primeiramente por escolher me orientar e assim conduzir uma orientação firme e de excelência. Agradeço imensamente a disponibilidade constante, inclusive em finais de semana e feriados, as oportunidades oferecidas e a confiança em mim depositada. Agradeço as palavras duras nos momentos pertinentes, pois o crescimento é um processo de erros e acertos mas sobretudo de coragem e discernimento. Toda minha gratidão por mostrar que deve-se buscar sempre além. Obrigada pela inspiração nesses dois anos tão importantes na minha formação profissional.

À Prof<sup>a</sup>. Dr<sup>a</sup>. Cristiane Ribeiro Salmon, pela co-orientação e todos os ensinamentos transmitidos desde o início do projeto, pela paciência e acolhimento, ao suporte inestimável na

fase de experimento animal, além da presença nas demais fases de experimento. Obrigada pela condução, por ensinar o caminho e acreditar que os passos seriam dados.

À Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), pela bolsa concedida durante o mestrado (2015-2017) e à Fundação de Amparo a Pesquisa no Estado de São Paulo (FAPESP) pelo auxílio (2015/06372-2) de pesquisa concedido.

Aos professores do Programa de Pós-graduação em Clínica Odontológica, área de concentração Periodontia, Prof. Dr. Antonio Wilson Sallum, Prof<sup>a</sup>. Dr<sup>a</sup>. Denise Carleto Andia, Prof. Dr. Enilson Antonio Sallum, Prof. Dr. Francisco Humberto Nociti Júnior, Prof<sup>a</sup>. Dr<sup>a</sup>. Karina Gonzales Silvério Ruiz, Prof. Dr. Marcio Zaffalon e Prof. Dr. Renato Corrêa Viana Casarin, pelas disciplinas ministradas durante o mestrado, assim como os ensinamentos transmitidos e convivência na clínica, que foram fundamentais no processo ensino-aprendizagem.

Aos professores da banca de qualificação: Prof. Dr. Renato Corrêa Viana Casarin, Prof<sup>a</sup>. Dr<sup>a</sup>. Denise Carleto Andia, Prof<sup>a</sup>. Dr<sup>a</sup>. Samira Salmeron, Prof<sup>a</sup>. Dr<sup>a</sup>. Luciane Martins pelo aceite do convite e pela criteriosa avaliação do trabalho, contribuindo de forma considerável com o aprimoramento do trabalho. Aos professores titulares e suplentes da banca de defesa da dissertação do mestrado, Prof. Dr. Francisco Humberto Nociti Júnior, Prof<sup>a</sup>. Dr<sup>a</sup>. Bruna Rabelo Amorim, Prof<sup>a</sup>. Dr<sup>a</sup>. Karina Gonzales Silvério, Prof<sup>a</sup>. Dr<sup>a</sup>. Daiane Cristina Peruzzo e Prof<sup>a</sup>. Dr<sup>a</sup>. Samira Salmeron, por aceitar compor a banca e disponibilizar tempo e atenção para o aperfeiçoamento do trabalho apresentado.

Ao Prof. Dr. Prof. Dr. Pedro Duarte Novaes pela solicitude e gentileza durante as fases do trabalho realizadas no departamento de Morfologia, área Histologia e Embriologia, agradeço ainda à equipe técnica nas pessoas de Adriano Luis Martins, pela paciência e ensinamentos sobre as diversas abordagens em microscopia, e Flávia Sammartino Mariano Rodrigues, pela extrema prestimosidade e delicadeza na condução dos trabalhos. À Gabriel Abuna (Materias Dentários), pelo aporte, obsequiosamente compartilhando os conhecimentos frente à microscopia confocal.

À equipe técnica da Coordenadoria de Pós-Graduação, em especial à Ana Paula Carone pela afabilidade e presteza na realização da sua função. Agradeço à Regina Célia Corrêa Caetano da Silva, da secretaria da área de Periodontia, pela extrema competência profissional e, sobretudo, pelo ser humano sensacional que por diversos momentos ofereceu amparo. À Eliete Ferreira Lima Marim pela eficiência e o sorriso acolhedor. À Mariana Piovezan Fugolin pelo suporte no laboratório de Periodontia e por tornar o ambiente mais aprazível.

Agradeço aos amigos e colegas da área de Periodontia: Amanda Bandeira, Ana Livia Fileto, Fernanda Felix, Guilherme de Oliveira, Isabela Lima França, João Paulo Sangiorgio, Manuela Rocha, Marcela Di Moura, Maria Alice Gatti Palma, Mayra Albiero, Mabelle Monteiro, Mércia Cunha, Miki Saito, Rafaela Videira, Rahyza Freire, Tiago Ozi, Tiago Taiete, Tiago Tarbes e Viviene Barbosa pela agradável convivência diária e pelo espírito de cooperação.

À Amanda Bandeira, pela amizade e companheirismo. Pela presença constante em todos os momentos, por ser esse ser único em espontaneidade, que torna os dias mais leves. À Rafaela Videira, por ser essa pessoa apaixonada, que trabalha com afinco, obrigada pela amizade e tranquilidade inerentes à sua essência. À Rahyza Freire, pessoa na qual descobri muitas afinidades e se tornou o ombro fraterno, a nitidez nos momentos mais turvos. Obrigada pelas palavras sensatas e parceria “deboísta”. Tenho certeza que a amizade deste grupo transcenderá a fase de pós-graduação.

À Mércia Cunha e Viviene Barbosa, pelo acolhimento inicial e todo suporte no período de experimentos animais no biotério. Além dos momentos agradáveis que compartilhamos dentro e fora da faculdade.

À Samira Salmeron, pelo carinho e atenção. Agradeço o suporte no biotério e os ensinamentos profissionais e pessoais nos períodos de permanência na sala de estudo da Periodontia.

À Ana Livia Fileto, Fernanda Felix, Manuela Rocha e Mabelle Monteiro, pela recepção afetuosa. O cuidado constante e atenção com os recém-chegados foi primordial para atravessar esses dois anos.

À Marcela Di Moura, por ser luz, por ser o sorriso largo nas horas mais difíceis, por ser um ser humano único que ensina o que sabe, faz sem esperar nada em troca, trabalha com amor e alegria, alguém que sei que posso contar a qualquer hora.

Aos amigos graduados na Universidade Federal da Paraíba e também integrantes do Programa de Pós-Graduação da FOP/UNICAMP: Ana Camila Assis (Cariologia), Bruno Mariz (Patologia), Diego Nóbrega (Cariologia), Emerson Tavares (Pediatria), Irlan Freires (Farmacologia, Anestesiologia e Terapêutica), Jaíza Araújo (Farmacologia, Anestesiologia e Terapêutica), Jossaria Pereira (Pediatria), Laíse Lima (Anatomia), Livia Alves (Microbiologia e Imunologia), Louise Dornelas (Prótese), Marina Moreno (Materiais dentários), Mayara Abreu (Prótese), Renally Wanderley (Materiais dentários) e Rodrigo Lins (Dentística) pelos momentos agradáveis, por tornar a FOP um lugar mais familiar.



À Isaac Jordão, pelo incentivo em ingressar no Programa de pós-graduação da FOP e pelo afável suporte na fase de adaptação.

À Renally Wanderley, pela acolhida. À Marina Moreno, pelos conselhos, ensinamentos, pelo cuidado, por ser o sorriso amigo nos corredores da Faculdade.

Aos amigos da área de Pediatria: Cláudia Lobelli, Filipe Martins e Samuel Chaves por tornar a FOP um lugar mais leve e iluminado. À Manoelito Silva Júnior (Saúde Coletiva), pelos momentos agradáveis e amizade, além das discussões científicas e de vida.

À Louise Moraes Dornelas Bezerra, dupla na Graduação, companheira de viagens, minha procuradora, amparo na pós-graduação, família em Piracicaba. Agradeço pelo suporte inestimável nestes mais de 7 anos de amizade.

À Rodrigo Barros Esteves Lins, amigo da Graduação, por também ser família em Piracicaba, pela companhia nesta jornada. Obrigada pelo ouvido amigo, pela generosidade nos atos, pela integridade inspiradora e por me fazer uma pessoa melhor.

Aos queridos amigos do departamento de Prótese e Periodontia, área prótese: Bruna Ximenes, Camilla Fraga, Heloísa Pantaroto, Jairo Cordeiro, Louise Dornelas, Mariana Barbosa, Mayara Abreu e Olívia Figueredo por serem as melhores pessoas para conviver, pela troca de experiências, o cuidado com o outro e estímulo acadêmico.

Aos funcionários do Biotério, pela paciência e colaboração. À equipe da biblioteca da FOP. Aos funcionários da limpeza, manutenção e segurança que prestam serviço à FOP/UNICAMP, pela colaboração sempre que solicitada, pelas saudações matinais com sorriso no rosto de todos os dias, pelo zelo conosco aos finais de semana e horários menos convencionais.

## EPÍGRAFE

***“Não enterrem os talentos! Apostem em ideais grandes que alargam o coração, ideais de serviço que tornam fecundos os seus talentos. A vida não nos foi dada para que a conservemos para nós mesmos, mas nos foi dada para que a doemos. Caros jovens, tenham uma grande coragem! Não tenham medo de sonhar coisas grandes!”***

Papa Francisco

***“A dor é inevitável, mas o sofrimento é opcional”***

Sri Sri Ravi Shankar

## RESUMO

Embora tenha havido avanço no entendimento da homeostase do cimento dental, o papel dos cementócitos permanece obscuro. Este estudo buscou caracterizar morfológica e ultraestruturalmente eventos relevantes nos componentes celulares do cimento dental durante a formação de novo cimento. Foi utilizado o modelo de indução de aposição do cimento dental nos molares inferiores de camundongos Swiss Webster por extração dos seus antagonistas. Para análise através de microscopia confocal os animais foram separados em um grupo induzido e um controle, e sacrificados após períodos de 6 e 21 dias ( $n=5$ ). Foi mensurada a área de cimento dental na região apical dos grupos, além da quantidade de células nesta área delimitada, a relação célula/área e área do corpo dos cementócitos. Ademais, foi realizada uma análise morfológica da rede de canalículos. Posteriormente, animais do grupo induzido e controle foram eutanasiados após 21 dias de experimento ( $n=3$ ), os cortes foram preparados e analisados por microscopia eletrônica de varredura (MEV). Finalmente, após 21 dias de experimento os animais ( $n=6$ ) foram perfundidos com solução de Karnovsky e as mandíbulas foram processadas para análise por microscopia eletrônica de transmissão (MET). Os dados mostraram aumento na área do cimento dental no grupo induzido, comparado ao controle aos 21 dias ( $p<0.05$ ), o que confirma a eficiência do modelo experimental na estimulação da aposição do cimento dental. No entanto, o número de cementócitos não foi alterado pela aposição de cimento, aos 21 dias foi identificada menor densidade celular na região estimulada ( $p<0.05$ ). A área do corpo celular dos cementócitos foi significativamente aumentada frente à aposição aos 21 dias ( $p<0.05$ ). As análises por microscopia eletrônica de transmissão demonstraram que, no grupo induzido, os cementócitos apresentaram corpos celulares e núcleos aumentados, além de maior quantidade de cromatina condensada na periferia e eucromatina espalhada no núcleo comparado ao controle. As análises por MEV revelaram dois padrões de lacunas no cimento celular: lacunas vazias e lacunas com remanescente celular, distribuídas irregularmente na matriz. Não foram observadas alterações organizacionais das fibras de Sharpey e dos feixes de fibras presentes na matriz. Desta forma, os achados morfológicos e ultraestruturais sugerem que os cementócitos, juntamente com cementoblastos, podem desempenhar um papel importante no controle da aposição experimental da matriz do cimento dental formado.

**PALAVRAS-CHAVE:** Cimento Dentário. Cementogênese. Microscopia.

## ABSTRACT

Despite significant advance in understanding dental cementum homeostasis, the role cementocyte remains unclear. This study aimed at describing morphologically and ultrastructurally relevant events in extracellular matrix and cellular components in the course of new cementum development. Experimentally-induced dental cementum apposition model was used. Extraction of the maxillary molars of Swiss Webster was performed inducing mandibular molar supereruption. A morphological analysis of canaliculi network and cementocytes was performed in decalcified sections with confocal microscopy. Animals were assigned into two groups and sacrificed after 6 and 21 days ( $n = 5$ ). Apical dental cementum area was measured, in addition the number of cells in this delimited area, the cell/ area ratio and body volume of the cementocytes. Sections on glass slides  $20\text{ }\mu\text{m}$  ( $n = 3$ ) were dehydrated, metallized and analyzed by scanning electron microscopy (SEM). Finally, after 21 days the animals ( $n = 6$ ) were perfused with Karnovsky's solution and mandibles processed for transmission electron microscopy (MET) analysis. Data demonstrated increased cementum area in experimental group compared to control at 21 days ( $p < 0.05$ ) confirming experimental model efficiency regarding dental cementum apposition. However, cementocyte number has not been changed by cementum apposition, at 21 days a lower cell density was found in the area where DC apposition was induced ( $p < 0.05$ ). Cementocyte volume was significantly increased when DC apposition was induced at 21 days ( $p < 0.05$ ). TEM analysis showed that, for the EIA group, cementocytes presented larger body and nuclei sizes, more peripherally condensed chromatin and euchromatin spread in the nuclei than the controls. Furthermore, SEM analysis revealed that the two patterns observed at the cellular cementum were either lacunae with cell remnants or empty lacunae, with the present cementocytes occupying lacunae irregularly shaped and unevenly distributed into the matrix. Moreover, Sharpey's and matrix fiber bundles were well organized with no evidence of clear organizational changes between the groups (control versus EIA). Therefore, morphological and ultrastructural data suggest that cementocytes, together with cementoblasts, may play an important role in controlling experimental dental matrix apposition.

**KEY WORDS:** Dental cementum. Cementogenesis. Microscopy.

SUMÁRIO

1 INTRODUÇÃO .....14

2 ARTIGO.....17

ULTRASTRUCTURAL ANALYSIS OF CEMENTOCYTES UNDER EXPERIMENTALLY-INDUCED DENTAL CEMENTUM APPOSITION .....17

3 CONCLUSÃO.....47

REFERÊNCIAS .....48

ANEXO 1. Certificado de aprovação do Comitê de Ética no Uso de Animais – Universidade Estadual de Campinas/UNICAMP.....52

ANEXO 2. Comprovante de submissão do artigo.....53

## 1 INTRODUÇÃO

O cimento dental é um tecido mineralizado que cobre a superfície radicular do dente e promove por meio do ligamento periodontal, a inserção deste no osso alveolar (Bosshardt & Selvig, 1997; Foster & Somerman, 2012). Este tecido é composto por dois tipos celulares, os cementoblastos e os cementócitos, que tem origem nas células mesenquimais indiferenciadas do tecido conjuntivo do folículo dental durante a odontogênese (Bosshardt & Selvig, 1997). Os cementoblastos produzem a matriz extracelular por aposição de camadas. Durante o processo de cementogênese, enquanto muitos cementoblastos se alinham formando uma camada celular ao longo da superfície do cimento, alguns cementoblastos são aprisionados na matriz recentemente produzida e tornam-se cementócitos, a semelhança do tecido ósseo onde osteoblastos tornam-se osteócitos aprisionados. Da mesma forma que os osteócitos na matriz mineralizada do tecido ósseo, os cementócitos residem em lacunas no interior da matriz mineralizada do cimento dental e comunicam-se entre si e com os cementoblastos localizados na superfície do cimento por meio de prolongamentos citoplasmáticos que se estendem dentro de canalículos e canais (Bosshardt & Selvig, 1997; Bosshardt & Schroeder, 1996; Saygin, Giannobile & Somerman, 2000) .

Por muitos anos, o papel dos osteócitos na manutenção da homeostasia do tecido ósseo foi subestimado ou até mesmo ignorado, uma vez que se acreditava que estas eram células inativas. Mais recentemente, vários grupos têm trabalhado no sentido de desvendar a real contribuição dos osteócitos na manutenção do equilíbrio do tecido ósseo (Dallas, Prideaux & Bonewald, 2013; Bellido, 2014). Trabalhos recentes, in vivo e in vitro, desvendaram muitas das funções suspeitas dos osteócitos, como mecano-sensorial e regulação da remodelação óssea e mineralização (Zhao et al., 2002; Feng et al., 2006; Gluhak-Heinrich et al., 2003; Kondoh et al., 2014; Lau et al., 2013). Além disso, estudos avançam em funções inovadoras dos osteócitos, tais como a capacidade de agir como um órgão endócrino, sugerindo um potencial para influenciar células e tecidos muito além do microambiente ósseo (Dallas, Prideaux & Bonewald, 2013).

Alguns estudos sugerem que a sinalização entre osteócitos e a paratireoide e rim indica a importância da função endócrina do osteócitos e o impacto que a manutenção da viabilidade dos osteócitos pode ter em outros tecidos (St John et al., 2014; Ito et al., 2015; Lindberg et al., 2014).

Neste contexto, o entendimento da participação de importantes vias de sinalização, como a via Wnt, e da secreção de moléculas chave no processo de remodelação do osso como a esclerostina (Compton & Lee, 2014; Honasoge, Rao & Rao, 2014; Sapir-Koren & Livshits, 2014) e o fator de crescimento 23 de fibroblastos (FGF23) (Lindberg et al., 2014; Sapir-Koren & Livshits, 2014) e da proteína de matriz dentinária (DMP1) (Feng et al., 2003; Toyosawa et al., 2012; Lee, Yamaguchi & Iimura, 2014) principalmente por osteócitos, foram os avanços recentes mais relevantes na biologia do esqueleto.

Diferente do tecido ósseo, o cemento não possui um papel de “*turnover*” ou remodelação significativos, ou seja, é capaz de reparar-se a um grau limitado e não é reabsorvido em condições normais (Bosshardt & Selvig, 1997; Foster & Somerman, 2012). Além disso, o cemento responde diferentemente a intervenções terapêuticas nos casos em que os tecidos periodontais são perdidos como consequência da doença periodontal, onde a regeneração do cemento é frequentemente difícil e imprevisível (Bosshardt & Sculean, 2009). Estudos têm sido conduzidos no sentido de tentar identificar fatores específicos do cemento dental (Foster et al., 2007; Foster et al., 2011; Foster et al., 2012; Salmon et al., 2013; Nociti & Somerman, 2013) e na compreensão dos seus processos fisiológicos que possam contribuir com o desenvolvimento de técnicas regenerativas mais confiáveis e previsíveis. Entretanto, o conhecimento a nível celular e molecular do cemento dental, tanto durante a cementogênese ou como no tecido maduro em condições fisiológicas normais ou alteradas ainda está pobremente definido (Saygin, Giannobile & Somerman, 2000). Evidências recentes sugerem que cementócitos não apenas dividem características morfológicas com osteócitos, mas também expressam inúmeros reguladores do metabolismo de tecidos mineralizados em comum. Foi demonstrado que, similarmente aos osteócitos, cementócitos expressam a esclerostina (Jager et al, 2010; Lehen et al., 2012; Kuchler et al., 2014) e em estudo com animais “*knockout*” ambos os fenótipos de osso e

cimento foram alterados (Kuchler et al., 2014). Ainda, paralelamente ao que ocorre com osteócitos, importantes reguladores do metabolismo de tecidos mineralizados, incluindo DMP1, são expressos por cementócitos e podem ser regulados via hormônio 1,25(OH)2D3 (1,25D) (Nociti et al., 2014) e Fgf23 (Chu et al., 2010). Assim como osteócitos, cementócitos expressam fatores importantes na regulação do metabolismo do fosfato (Pi), incluindo proteína de anquilose progressiva (ANK), e animais “knockout” para essa proteína desenvolvem fenótipos alterados em tecidos ósseos e dentais (Foster et al., 2011; Nociti et al., 2002).

Embora seja sugerido que cementoblastos, juntamente com células do ligamento periodontal e do osso alveolar, são essenciais para a manutenção da homeostasia dos tecidos periodontais (Saygin, Giannobile & Somerman, 2000; Hefti, 1993; Bartold, 1995; Grzesik & Narayanan, 2002), o papel dos cementócitos nesse processo permanece completamente desconhecido. Dadas as características morfológicas e biológicas comuns para osteócitos e cementócitos, além das similaridades entre a matriz do tecido ósseo e do cimento dental, nossa hipótese é de que cementócitos possam exercer um papel significativo para a manutenção não apenas do cimento dental, mas também de todo o conjunto que compõe o periodonto de sustentação. E apesar de anos de pesquisa e desenvolvimento de conhecimento sobre o papel do cimento, o qual acredita-se ser importante no processo reparativo após a doença periodontal, ainda há aspectos relevantes em relação a este tecido que permanecem não esclarecidos.

Desta forma, o presente estudo foi realizado utilizando-se técnicas celulares de alta resolução para fornecer informações sobre a morfologia e ultraestrutura dos cementócitos, com o objetivo de determinar o papel exercido pelos cementócitos na homeostasia dos tecidos periodontais. O melhor conhecimento sobre o perfil celular dos cementócitos poderá fornecer novos “insights” sobre o tecido que possam permitir o desenvolvimento de terapias eficientes e mais previsíveis para regeneração dos tecidos periodontais.



**2 ARTIGO****ULTRASTRUCTURAL ANALYSIS OF CEMENTOCYTES UNDER  
EXPERIMENTALLY-INDUCED DENTAL CEMENTUM APPPOSITION**

Elis Janaina Lira dos Santos<sup>1</sup>, Cristiane Ribeiro Salmon<sup>1</sup>, Amanda Bandeira de Almeida<sup>1</sup>, Enilson A. Sallum<sup>1</sup>, Marcio Z. Casati<sup>1</sup>, Karina G. Ruiz<sup>1</sup>, Renato Casarin<sup>1</sup>, Kamila Rosamilia Kantovitz<sup>2</sup>, Francisco Humberto Nociti Júnior<sup>1,\*</sup>

<sup>1</sup> Department of Prosthodontics and Periodontics, Division of Periodontics, Piracicaba Dental School, State University of Campinas, São Paulo, Brazil.

<sup>2</sup> Department of Dental Materials, São Leopoldo Mandic Research Center, Campinas, São Paulo, Brazil

Corresponding author:

Francisco Humberto Nociti Júnior

Avenida Limeira, 901, Bairro Areião, Piracicaba, São Paulo 13414-903, Brazil.

Telephone: +55-19-21065301; Fax: +55-19-21065301

[nociti@unicamp.br](mailto:nociti@unicamp.br)

## ABSTRACT

**Background:** Cementocytes not only share morphological features with osteocytes, but also express several common biomarkers. Nevertheless, in contrast to osteocytes, there is a lack of evidence on the potential role of cementocytes in tissue homeostasis. Here, it is hypothesized that cementocytes are key players in dental cementum biology. **Material and Methods:** This study used a model of experimentally induced dental cementum apposition (EIA) in mice to determine whether cementocytes are responsive cells in such environment. Mandibular 1<sup>st</sup> molars were randomly induced to erupt for 6 and 21 days after extracting opposing maxillary molars, whereas contralateral teeth were used as controls. Histological sections were prepared from the 1<sup>st</sup> molars' region for transmission and scanning electron microscopy (TEM and SEM, respectively), whereas non-decalcified sections were labeled with fluorescein isothiocyanate and observed under a confocal microscope. Quantitative data was submitted to two-way ANOVA ( $\alpha=5\%$ ) followed by Tukey's test ( $\alpha=5\%$ ). **Results:** Data analysis showed an increased area of dental cementum (DC) for the EIA group compared to control at 21 days ( $p<0.05$ ), and therefore, confirmed DC apposition. Intriguingly, as cementocyte's number was not affected by DC apposition, at 21 days a lower cell density was found in the area where DC apposition was induced ( $p<0.05$ ). Cementocyte volume was significantly increased when DC apposition was induced at 21 days ( $p<0.05$ ). TEM analysis showed that, for the EIA group, cementocytes presented larger body and nuclei sizes, more peripherally condensed chromatin and euchromatin spread in the nuclei than the controls. Furthermore, SEM analysis revealed that the two patterns observed at the cellular cementum were either lacunae with cell remnants or empty lacunae, with the present cementocytes occupying lacunae irregularly shaped and unevenly distributed into the matrix. Moreover, Sharpey's and matrix fiber bundles were well organized with no evidence of clear organizational changes between the groups (control versus EIA). **Conclusion:** Together, these findings provide new insights on DC biology and reveal cementocytes as potential targets for tissue engineering applications.

**KEY WORDS:** dental cementum, cementogenesis, ultrastructure

## INTRODUCTION

Dental cementum (DC) is a “bonelike” mineralized connective tissue covering the root surface of teeth serving as the attachment structure for periodontal ligament (1). Both, DC and bone, present a highly similar structure and composition with a comparable mineral-to-organic matrix ratio (2). Nonetheless, mature bone has a lamellar organization and actively participates in the metabolism of the body, providing a reservoir for calcium and other elements (3-6); whereas DC is not innervated, avascular, less cellular and exhibits little or no remodeling, but continues to grow in thickness throughout life (1, 7-9). DC is composed of two cells types, cementoblasts and cementocytes, both originated from mesenchymal stem cells during odontogenesis. Cementoblasts have been described as large and cuboidal cells with euchromatin-rich nuclei, plentiful rough endoplasmatic reticulum (RER) and Golgi apparatus, whereas cementocytes reside in lacunae and exhibit canalicular network and irregular spacing (10).

The hallmarks of periodontal disease are the destruction of soft connective tissues, bone loss, and loss of connective tissue attachment to DC; these alterations, if left untreated, may lead to tooth loss (11-13). The aim of periodontal therapy is to regenerate and restore the various periodontal components affected by disease. New therapeutic approaches available include the use of barrier membranes for guided tissue regeneration, and applying growth factors and enamel matrix proteins to root surfaces (14-16); however, the effectiveness of these approaches is not predictable, especially when considering new cementum and attachment formation. A significant barrier to improved therapies for regeneration is that cementum biology remains poorly understood. Fundamental questions centered on cementocytes include whether they are physiologically active cells, whether they remain in communication with cells in the periodontal ligament (PDL) region, whether they are involved in cementum formation and repair, including response of cementum to altered demands of tooth use, and whether they regulate osteoclast activity at the cementum surface. Recent studies suggest that cementocytes, apparently terminally differentiated cells, present an *in vivo* expression profile that parallels that of osteocytes, including the expression of dentin matrix protein 1 (DMP1), sclerostin (SOST), E11/gp38, tumor

necrosis factor receptor superfamily, member 11b (osteoprotegerin/OPG), and tumor necrosis factor superfamily member 11 (RANKL) (17-20). In addition, using the experimentally-induced apposition (EIA) model to induce DC apposition, followed by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS) proteomic analysis, our group provided initial evidence that cementocytes may have an active role in neocementogenesis (21). Although, some progress has been made in terms of improving comprehension on the biology of DC, it is presently unknown whether cementocytes have *in vivo* functions related to cementum formation or repair. Here, it was asked whether cementocytes are dynamic actors in neocementogenesis capable of undergoing morphological and ultrastructural changes as a response to the continuous apposition of DC in a mouse model.

## MATERIALS AND METHODS

**Animals and surgeries:** Animal procedures were performed at the Piracicaba Dental School Animal Facility in accordance with the guidelines of the University Committee for Ethics in Animal Research (Protocol#3788-1). The experimentally-induced apposition model (EIA) was used to promote dental cementum (DC) apposition, as previously proposed (22, 23). Briefly, male Swiss Webster mice (35 days old) were sedated and the right/left maxillary molars were randomly extracted to allow DC apposition on the antagonist mandibular 1<sup>st</sup> molars. The contralateral side was used as the control group. After 6 and 21 days (21) of DC apposition, 19 animals were euthanized by cervical dislocation, the mandibles dissected and placed either in paraformaldehyde 4% for confocal imaging, formalin 4% for scanning electron microscopy (SEM) and Karnovsky's solution for transmission electron microscopy (TEM).

**Fluorescence analysis by confocal microscopy:** In order to access changes in cementocyte morphology as a consequence of the process of continuous neocementogenesis, undecalcified samples were labeled with fluorescein isothiocyanate (FITC) and analyzed by confocal microscopy. Experimentally-induced DC apposition was performed and the animals euthanized after 6 and 21 days (n=5). After euthanize, mandibles were dissected, fixed in 4% paraformaldehyde at 4°C under agitation for 24 hours, dehydrated in 70% ethanol for 24 hours, and embedded in SamplKwick Fast Cure Acrylic resin (Buehler, Lake Bluff, Illinois, USA), following the manufacturer's instructions. A diamond low-speed saw (South Bay Technology, San Clemente, California, USA) was used to obtain 400-500 µm bucco-lingual sections from the mesial root of the 1<sup>st</sup> mandibular molar, and then manually ground to a final thickness of approximately 300 µm. Sections were dehydrated in graded ethanol series (50%, 70%, 90% for 10 minutes each, and 100% for 30 minutes), and stained with 1% fluorescein isothiocyanate (FITC) for 4 hours at room temperature protected from light. Next, samples were washed in 100% ethanol for 30 minutes, let to dry at room temperature for 2 hours and mounted on Entellan®. The cellular cementum was

immediately analyzed in a confocal microscope with the laser set to 488 nm (Leica, Wetzlar, Hessen, Germany). Six serial images equally distant (0.8  $\mu\text{m}$ ), representative of the whole tooth, were obtained from Z-stack by confocal laser scanning microscopy (CLSM) in the bucco-lingual direction for five animals. Cellular cementum area, cementocyte number and density were determined, by a blind and calibrated examiner (EJLS), on the mesial root of the mandibular 1<sup>st</sup> molar for both, control and EIA groups, using an image-analysis system (Image J – Java-based image processing software package v3.91, National Institutes of Health (NIH), Bethesda, Maryland, USA). In addition, dimensional changes were accessed in twelve randomly selected cementocytes according to its location: close to dentin (n=6) and close to the periodontal ligament (n=6). Histological measurements were calculated as an average value for each animal and a final average determined for each experimental group. Figure 1 schematically illustrates all the accessed parameters.

**Scanning Electron Microscopy (SEM):** SEM analyses were performed on hemi-mandibles from mice under EIA and controls (n=3). At 21 days, animals were sacrificed, mandibles were dissected, fixed in 4% neutral formalin for 48 h and subsequently demineralized in 10% EDTA solution at 4°C under agitation for 17 days. Bucco-lingual paraffin serial sections (20  $\mu\text{m}$ ) were obtained from the mesial root of the 1<sup>st</sup> mandibular molar. Next, sections were dehydrated (3x, xylene), air-dried for 10 minutes at room temperature, dried to the critical point and gold coated (Denton Vacuum Desk II, Moorestown, New Jersey, USA). Images were obtained and analyzed in a scanning electron microscope (JEOL JSM 5600LV, Tokyo, Japan) operating at 15 kV in electron mode. A blinded examiner (EJLS) performed all the SEM analysis.

**Transmission Electron Microscopy (TEM):** In order to perform TEM analysis, animals under EIA and control (n=6) were perfused with Karnovsky's fixative solution (4% paraformaldehyde, 2% glutaraldehyde in Sorensen phosphate buffer 0.1M, pH 7.4) at 21 days after EIA. Mandibles were dissected and post-fixed in Karnovsky's solution for 24 hours, washed in PBS, decalcified in 10% EDTA for 17 days. Next, samples were

washed in PBS, post-fixed in 1% osmium tetroxide ( $\text{OsO}_4$ ), washed in PBS, dehydrated in a graded acetone series (75%, 95%, 100%) and embedded in LR White resin (EMS, Hatfield, Pennsylvania, USA). Thin sections (70-90 nm) were obtained from mesial root of 1<sup>st</sup> mandibular molars in a bucco-lingual direction, transferred to carbon-coated. Formvar grids, stained with uranyl acetate (10 minutes) followed by lead citrate (2 minutes). Mesh grids were then viewed and imaged on a transmission electron microscope JEM 1400 (Jeol, Japan). A blinded examiner (EJLS) performed all the TEM analysis.

**Statistical analysis:** DC area, cell number, cell density and dimensional changes were statistically compared between EIA and control groups using the two-way ANOVA followed by the Tukey's test ( $\alpha=0.05$ ) and Shapiro–Wilk test (SAS system for Windows v9.4 - SAS Institute, Cary, NC, USA).

## RESULTS

**Confocal analysis:** CLSM allowed qualitative and quantitative assessments. Qualitative analysis focused on the cementocytes' canalicular system, whereas quantitative parameters included the impact of EIA on cellular cementum area, cementocyte number, cellular cementum density (*e.g.*, number of cementocyte cellular / cellular cementum area) and cementocyte body size (figure 1). Data analysis showed that the cellular cementum area was increased by EIA at 21 days ( $p<0.05$ ), whereas at 6 days, no clear effect of EIA was found on the area of cellular cementum ( $p>0.05$ ) (Figures 2A-C). Intragroup analysis revealed that the number of cementocytes was increased over time for both groups ( $p<0.05$ ); however, intergroup comparisons showed that there was no difference between the experimental groups at 6 and 21 days (Figure 3A). Therefore, as cellular cementum area was significantly increased by EIA at 21 days and cementocytes number was not affected, cellular density was significantly decreased by EIA at 21 days (Figure 3B). Since cellular dimensional changes have been reported as an important characteristic of metabolic active cells, we examined whether

cementocytes' body size was affected in response to an experimental neocementogenesis model (EIA). Intragroup analysis showed that cementocytes' body size was significantly increased in response to continuous neocementogenesis ( $p < 0.05$ ), whereas in the control group, cementocytes' body size decreased overtime ( $p < 0.05$ ). Furthermore, at 21 days, intergroup analysis demonstrated that EIA resulted in larger cementocytes as compared to the control group (Figure 4A-B). In the current study, we found no relevant qualitative differences either when control and EIA groups were compared or when cementocytes and osteocytes were comparatively analyzed (Figure 5).

**Scanning Electron Microscopy (SEM) findings:** In the present investigation, SEM analysis was also used to qualitatively examine the impact of EIA on cementocytes at 21 days. Overall, for both experimental groups, dental cementum consisted of masses of radially arranged, extrinsic Sharpey's fibers having surfaces covered by numerous intrinsic collagen fibers. These Sharpey's fibers maintained a perpendicular orientation to the cementum surface and emerged with the fibers of the periodontal ligament. Sharpey's and matrix fiber bundles were well organized with no evidence of clear organizational changes between the groups (control versus EIA). Moreover, SEM analysis revealed that the two patterns observed at the cellular cementum were either lacunae with cell remnants or empty lacunae, with the present cementocytes occupying lacunae irregularly shaped and unevenly distributed into the matrix. Figure 6 illustrates the representative SEM findings.

**Transmission Electron Microscopy (TEM) findings:** In the present study, TEM analysis was employed to explore potential changes on the cellular ultrastructure of cementocytes in response to the process of neocementogenesis. We found that cementocytes from both, control and EIA groups presented defined nuclei, intact nuclear membrane and relatively poorly developed rough endoplasmic reticulum (RER). Cementocytes from the EIA group featured increased cell and nuclei volume, more peripherally condensed chromatin and also and euchromatin spread in the nuclei



than the controls. Surprisingly, for the EIA experimental group, there was a trend towards a larger empty space between the cementocyte and its lacunae.

In the current investigation, we used this system to also direct our analyses to the cementoblasts. In general, for the EIA group, cementoblasts were identified as cells with abundant cytoplasm, nuclei with prominent nucleoli and rough endoplasmic reticulum (RER) with large number of minute granular particles (ribosomes) attached to the cisternae surface, which represents the typical cell morphology of high metabolic rate, indicating intense protein synthesis. In comparison with cementocytes, cementoblasts were larger cells presenting well-defined nuclei, intact nuclear membrane and well-developed RER (Figure 7).

## DISCUSSION

Dental cementum (DC), a mineralized tissue that covers the tooth root, is similar in several aspects to bone, however DC biology remains poorly understood. Previously thought to be passive cells residing within the bone matrix, currently osteocytes are accepted as active modulator of bone homeostasis and remodeling, and have been reported to actively contribute to endocrine regulation of mineral metabolism (4, 24). The panorama of what is known about DC comes from studies on its histology and composition (25, 26, 27), and although recent progress has been made (19, 21), there is still a lack of information related to the role of cementocytes in DC biology. Due to advances in the exploration of osteocytes as an important actor in bone metabolism (28, 29), comprehensive research on cementocytes has been encouraged in order to clarify the complete function of these cells in periodontal homeostasis and regeneration. Here, it was used the EIA model associated with *in vivo* fluorescent imaging, TEM and SEM analysis to test the hypothesis that cementocytes are active cells engaged with neocementogenesis. Our findings indeed demonstrate a number of evidences that cementocytes might react to continuous DC apposition, including increasing cementocytes' body size, increasing nuclei volume and chromatin

condensation distinctions. To the best of our knowledge, this is the first time that a potential role of cementocytes in neocementogenesis has been demonstrated.

Sicher and Weinmann's historic model of the unopposed rodent molar established the biological foundation for root cementum and bone growth as consequence of physiological movement. The authors suggested removing the upper molars on one side and allowing the lower molars to erupt without an antagonist. This model seems to be useful to study aspects of tissue remodeling during axial tooth movement (30, 31).

Authors have revealed that new cementum and alveolar bone are formed at the apex of the un-opposed molar after 12 days (32), therefore in the present study time points at 6 and 21 days were considered according to previous study using the same approach (21). In earlier periods initial molecular changes occur while in the late periods perhaps morphological changes have been occurred. Eruption and distal movement induced by unloading are result of distal bone resorption and mesial/apical bone apposition (22) that is the reason to evaluate mesial root in the present study.

A number of approaches have been used to determine whether distinct cell types are engaged with biological processes. As we have noted in previous ultrastructural analysis using TEM, cementoblasts are located nearby to external cementum layer and present evidences of metabolically active cells such as pronounced rough endoplasmic reticulum (RER). Cementocytes are irregular spacing in cellular cementum and differ widely in shape, some being elongated, more stellate or cuboidal, with relatively few organelles compared to cementoblasts (1, 27, 33). An important event in alveolar bone is the change from osteoblast to osteocyte, wherein cell morphology is substantially modified. Osteoblasts undergoes morphological change from polygonal to a more stellate shape with dendritic processes, once enclosed in bone tissue features a polarity to direct mineral formation (29). While osteocyte is embedded and differentiates, is observed cellular body contraction as well as reduction in the cytoplasmic organelles such as Golgi complex and endoplasmic reticulum (34, 35). Cementoblast to cementocytes transition demonstrate change in cellular shape and size, as well as decrease in cytoplasmic organelles, indicative of low metabolic rate (30). Indeed, in this study, experimental and control groups indicated some ultrastructural differences regardless of cementoblasts, which feature extensive

rough endoplasmic reticulum (RER) under stimuli, indicating increased cellular activity. Cementocytes also presented changes as larger cells and nuclei volume under experimentally-induced apposition, however RER are not extremely developed compared to cementoblasts. These findings may suggest that cementocytes does not produce protein similar to cementoblasts but may be participating in signaling for more active matrix apposition by cementoblasts.

Previous Scanning electron microscopy analysis was used to view transverse sections of mandibular 1<sup>st</sup> molar under EIA and revealed that the area of cellular cementum was increased at the root apex by 37% (32). In addition, our findings demonstrate that cellular cementum area was significantly increased by EIA at 21 days and cementocytes number was not affected. It was hypothesize, although cementocytes number has not been affected by EIA, cementocytes present at cellular cementum play an active role on neocementogenesis and these findings suggest EIA model affect cementocyte body size instead cementocyte number. Some authors have shown that increased cell volume is associated with cellular activity (36, 37). Overall, the ultrastructural results of this study suggest that experimentally-induced dental cementum apposition model may promote changes in the cellular machinery once the cellular volume is increased and this is not related to increase cementocytes in number.

Osteocyte reside within a ellipsoid and regular lacunae, their dendritic connections provide communication between osteocytes and surface cells like osteoblasts and osteoclasts as well as with distant cells via signaling (38, 39). These long cytoplasmic processes radiate in different directions surround cell body. This presents significant value to overcome the matrix enclosed environment and to establish communication with other cells as well as access nutrients and oxygen (4, 40). Analysis using SEM have demonstrated that similar to osteocytes, cementocytes occupy lacunae and display projections within a canalicular network, suggesting they are adapted as cells embedded within a extracellular mineralized matrix, establishing communication system with one another and with cells outside the cellular cementum (4, 10, 41).

Although studies have shown that osteocytes embedded in mineralized bone matrix play an important role in bone tissue homeostasis, the role of cementocytes in dental cementum has not been thoroughly elucidated in terms of associated cells and

regulatory factors involved in pathological processes and other conditions. Periodontal disease, for instance, refers to the inflammatory processes that occur in the tissues surrounding the teeth in response to periodontopathogenic microorganisms (42, 43), and neutrophils, macrophages and lymphocytes play important role in the host local response (44). Moreover, immunoinflammatory response includes events as activation of complement system, antigen presentation to T lymphocytes, antibodies production and cytokines and chemokines expression (45, 46, 47). Tumor necrosis factor (TNF- $\alpha$ ), Interleukin-1 $\beta$  (IL-1 $\beta$ ) and Interleukin-6 (IL-6) are the main cytokines expressed. TNF- $\alpha$  regulate adhesion molecules and stimulate the production of new chemokines, matrix metalloproteinases and RANKL. IL-1 $\beta$  and IL-6 are associated with migration of inflammatory cells and osteoclastogenesis (48, 49, 50). Moreover, osteoclastogenesis mechanism in periodontal disease is associated to these regulatory molecules. Osteoblasts express RANKL, the main regulator of osteoclast differentiation and activation, osteoclast cells induce bone resorption (51, 52). On the other hand, the inhibitor of osteoclastogenesis, osteoprotegerin (OPG) upon binding to the RANK receptor, present in the preosteoclasts, block bone resorption (53, 54). Recently, osteocytes have been considered the major source of RANKL and may have greater capacity to support osteoclastogenesis (55). Bone lacunar canalicular system seems to conduct bone remodeling and resorption, transmitting signals to coordinate osteoblast and osteoclast activity (56, 57). *In vivo* experimental model in mice for apical periodontitis demonstrated bone resorption and as the disease progresses RANKL/OPG expression has been increased. In addition, periodontal ligament and cementum reabsorption has been detected. Overall, the study suggest cementocytes express RANKL in response to endodontic infection and may be involved in the pathogenesis of apical periodontitis (20). DC and bone, present a highly similar structure and composition, the fact that cementocytes express RANKL may suggest these cells play an active role in modulate development of periodontal disease. Similar to osteocyte, cementocytes also express Sclerostin (SOST) (18), DMP1 (58) and important factors in the regulation of phosphate metabolism (Pi), including progressive ankylosis protein (ANK) (59, 60). Therefore, these findings strongly suggest that cementocytes are involved in events such as the development of resorption and

perhaps cell recruitment to inflammatory sites, due to their similarity to osteocytes. However, further studies have been encouraged to prove this important role.

A similar study using the same model of experimentally-induced apposition (EIA) in mice and the same time points (6 and 21 days) identified proteins associated with neocementogenesis. These encompassed SerpinF1, tenascin-X (TNX), aspirin (ASPN), collagen type XI alpha 1 (COL11A1), osteonectin (SPARC), thrombospondin 1 (THBS1), osteopontin (OPN), and fibromodulin (FMOD). Overall, proteomic analysis demonstrated significantly altered protein profile in dental cementum under EIA. The study demonstrated the model efficiency similar to our findings (21).

Recently, with the establishment of improved therapies for periodontal tissues, regenerative approach following periodontal disease it has become one of the main goals in periodontal therapy. Future studies focusing on defining the mechanisms involved in cementogenesis should assist in establishing the significance of cementocytes in modulating cementum formation and clarify their effects on the periodontium. Our findings indicate that cementum is responsive to experimentally-induced apposition model, due to an increased area of dental cementum (DC) after stimuli. Ultrastructural changes including increasing cementocytes' body size, increasing nuclei volume and chromatin condensation suggest that cementocytes may play active role in dental cementum homeostasis.

#### **CONFLICT OF INTEREST**

The authors report no conflict of interest.

#### **ACKNOWLEDGMENTS**

This work was supported by grant no. 2015/06372-2 from Agency São Paulo Research Foundation (FAPESP).

## REFERENCES

1. Bosshardt DD, Selvig KA. Dental cementum: the dynamic tissue covering of the root. *Periodontol 2000*. 1997;13:41-75.
2. Bosshardt DD. Are cementoblasts a subpopulation of osteoblasts or a unique phenotype? *J Dent Res*. 2005; 84:390–406.
3. Zhao S, Zhang YK, Harris S, Ahuja SS, Bonewald LF. MLO-Y4 osteocyte-like cells support osteoclast formation and activation. *J Bone Miner Res*. 2002;17:2068-79.
4. Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr Rev*. 2013;34:658-90.
5. Kondoh S, Inoue K, Igarashi K, Sugizaki H, Shirode-Fukuda Y, Inoue E, et al. Estrogen receptor alpha in osteocytes regulates trabecular bone formation in female mice. *Bone*. 2014; 60:68-77.
6. Kajiya H. Calcium signaling in osteoclast differentiation and bone resorption. *Adv Exp Med Biol*. 2012; 740:917-32.
7. Freeman E., 1994. *Oral Histology*, fourth ed. Mosby-Year Book Inc., St. Louis, MO, USA, pp. 276–312.
8. Grant DA, Stern IB, Listgarten MA, 1988. *Periodontics*, sixth ed. The CV Mosby Company, St. Louis, MO, USA, pp. 76–93.
9. Saygin NE, Giannobile WV, Somerman MJ. Molecular and cell biology of cementum. *Periodontol 2000*. 2000;24:73-98.

10. Zhao, N ; Foster, B.L ; Bonewald, L.F. The Cementocyte An Osteocyte Relative? Journal of Dental Research. 2016; 95(7), 734-741.
11. Page RC, Schroeder HE. Pathogenesis of inflammatory periodontal disease. A summary of current work. Lab Invest. 1976; 34(3): 235-49.
12. Martinez-Canut P. Predictors of tooth loss due to periodontal disease in patients following long-term periodontal maintenance. J Clin Periodontol. 2015; 42(12):1115-25.
13. Nibali L, Sun C, Akcalı A, Meng X, Tu YK, Donos N. A retrospective study on periodontal disease progression in private practice. J Clin Periodontol. 2016.
14. Sam G, Pillai BRM. Evolution of Barrier Membranes in Periodontal Regeneration-“Are the third Generation Membranes really here?”. J Clin Diagn Res. 2014 ; 8(12): ZE14–ZE17.
15. Darby I B , Morris K H. A Systematic Review of the Use of Growth Factors in Human Periodontal Regeneration. Journal of Periodontology. 2013; 84(4), 465-476.
16. Esposito M, Grusovin MG, Papanikolaou N, Coulthard P, Worthington HV. Enamel matrix derivative (Emdogain) for periodontal tissue regeneration in intrabony defects. A Cochrane systematic review. Eur J Oral Implantol. 2009; 2(4):247-66.
17. Ye L, Zhang S, Ke H, Bonewald LF, Feng J. Periodontal Breakdown in the Dmp1 Null Mouse Model of Hypophosphatemic Rickets. Journal of dental research. 2008;87(7):624-629.

18. Jager A, Gotz W, Lossdorfer S, Rath-Deschner B. Localization of SOST/sclerostin in cementocytes in vivo and in mineralizing periodontal ligament cells in vitro. *J Periodontal Res.* 2010;45:246-54.
19. Zhao N, Nociti FH, Duan P, Prideaux M, Zhao H, Foster BL, Somerman MJ, Bonewald LF. Isolation and Functional Analysis of an Immortalized Murine Cementocyte Cell Line, IDG-CM6. *J Bone Miner Res.* 2016.31: 430–442.
20. De Rossi A, Fukada SY, De Rossi M, da Silva RA, Queiroz AM, Nelson-Filho P, da Silva LA. Cementocytes Express Receptor Activator of the Nuclear Factor Kappa-B Ligand in Response to Endodontic Infection in Mice. *J Endod.* 2016 ;42(8):1251-7.
21. Salmon CR, Giorgetti AP, Paes Leme AF, Domingues RR, Sallum EA, Alves MC, Kolli TN, Foster BL, Nociti FH Jr. Global proteome profiling of dental cementum under experimentally-induced apposition. *J Proteomics.* 2016 Jun 1;141:12-23.
22. Luan X, Ito Y, Holliday S, Walker C, Daniel J, Galang TM, et al. Extracellular matrix-mediated tissue remodeling following axial movement of teeth. *J Histochem Cytochem.* 2007;55:127-40.
23. Walker CG, Dangaria S, Ito Y, Luan X, Diekwisch TG. 2010. Osteopontin is required for unloading-induced osteoclast recruitment and modulation of RANKL expression during tooth drift–associated bone remodeling, but not for super-eruption. *Bone.* 47(6):1020–1029.
24. Quarles LD. Endocrine functions of bone in mineral metabolism regulation. *The Journal of Clinical Investigation.* 2008;118(12):3820-3828.




25. Scivetti M, Pilolli GP, Corsalini M, Lucchese A, Favia G. Confocal laser scanning microscopy of human cementocytes: analysis of three-dimensional image reconstruction. *Ann Anat.* 2007; 189(2): 169-74.
26. Ho SP, Yu B, Yun W, Marshall GW, Ryder MI, Marshall SJ. Structure, chemical composition and mechanical properties of human and rat cementum and its interface with root dentin. *Acta biomaterialia.* 2009;5(2):707-718.
27. McKee MD, Zalzal S, Nanci A. Extracellular matrix in tooth cementum and mantle dentin: localization of osteopontin and other noncollagenous proteins, plasma proteins, and glycoconjugates by electron microscopy. *Anat Rec.* 1996; 245(2): 293-312.
28. Aarden EM, Burger EH, Nijweide PJ. Function of osteocytes in bone. *J Cell Biochem.* 1994;55(3):287-99.
29. Bonewald LF. The amazing osteocyte. *J Bone Miner Res.* 2011; 26(2): 229-38.
30. Sicher H, Weinmann JP. Bone growth and physiologic tooth movement. *Am J Orthod* 1944;30:109– 132.
31. Sicher H. Tooth eruption: Axial movement of teeth with limited growth. *J Dent Res* 1942;21: 395– 402.
32. Holliday S, Schneider B, Galang M T S, Fukui T, Yamane A, Luan X, Diekwisch TGH. BONES, TEETH AND GENES: A Genomic Homage to Harry Sicher's "Axial Movement of Teeth". *World J Orthod.* 2005; 6(1): 61–70.
33. Yamamoto T, Domon T, Takahashi S, Wakita M. 1996. Cellular cementogenesis in rat molars: the role of cementoblasts in the deposition of intrinsic matrix fibers of cementum proper. *Anat Embryol (Berl).* 193(5):495–500.

34. Palumbo C. A three-dimensional ultrastructural study of osteoid-osteocytes in the tibia of chick embryos. *Cell Tissue Res.* 1986; 246(1):125-31.
35. Dudley HR, Spiro D. The fine structure of bone cells. *J Biophys Biochem Cytol.* 1961;11:627–649.
36. Marguerat S. Coordinating genome expression with cell size. 2012; 28(11): 560–565.
37. Lang F, Föller M, Lang K, Lang P, Ritter M, Vereninov A, Szabo I, Huber SM, Gulbins E. Cell volume regulatory ion channels in cell proliferation and cell death. *Methods Enzymol.* 2007; 428:209-25.
38. Schaffler MB, Kennedy OD. Osteocyte Signaling in Bone. *Current Osteoporosis Reports.* 2012;10(2):118-125.
39. Webster DJ, Schneider P, Dallas SL, Müller R. Studying osteocytes within their environment. *Bone.* 2013;54(2):285-295.
40. Beno T, Yoon YJ, Cowin SC, Fritton SP. Estimation of bone permeability using accurate microstructural measurements. 2006; 39(13): 2378–2387.
41. Lanske B, Densmore MJ, Erben RG. Vitamin D endocrine system and osteocytes. *Bonekey Rep.* 2014; 5(3):494.
42. Socransky SS, Haffajee AD. The bacterial etiology of destructive periodontal disease: current concepts. *J Periodontol.* 1992;63(4):322-31.
43. Armitage GC. Clinical evaluation of periodontal diseases. *Periodontol* 2000. 1995 ;7:39-53.


44. Cekici A, Kantarci A, Hasturk H, Van Dyke TE. Inflammatory and immune pathways in the pathogenesis of periodontal disease. *Periodontology* 2000. 2014;64(1):57-80.
45. Hajishengallis G. Complement and Periodontitis. *Biochemical pharmacology*. 2010;80(12):1992-2001.
46. Gemmell E, Carter CL, Hart DN, Drysdale KE, Seymour GJ. Antigen-presenting cells in human periodontal disease tissues. *Oral Microbiol Immunol*. 2002;17(6):388-93.
47. Garlet GP, Martins W Jr, Ferreira BR, Milanezi CM, Silva JS. Patterns of chemokines and chemokine receptors expression in different forms of human periodontal disease. *J Periodontal Res*. 2003;38(2):210-7.
48. Kornman KS, Crane A, Wang HY, di Giovine FS, Newman MG, Pirk FW, Wilson TG Jr, Higginbottom FL, Duff GW. The interleukin-1 genotype as a severity factor in adult periodontal disease. *J Clin Periodontol*. 1997;24(1):72-7.
49. Irwin CR, Myrillas TT. The role of IL-6 in the pathogenesis of periodontal disease. *Oral Dis*. 1998;4(1):43-7.
50. Graves DT, Cochran D. The contribution of interleukin-1 and tumor necrosis factor to periodontal tissue destruction. *J Periodontol*. 2003;74(3):391-401.
51. Katagiri T, Takahashi N. Regulatory mechanisms of osteoblast and osteoclast differentiation. *Oral Dis*. 2002;8(3):147-59.
52. Atkins GJ, Kostakis P, Pan B, Farrugia A, Gronthos S, Evdokiou A, Harrison K, Findlay DM, Zannettino AC. RANKL expression is related to the differentiation state of human osteoblasts. *J Bone Miner Res*. 2003;18(6):1088-98.

53. Udagawa N, Takahashi N, Yasuda H, Mizuno A, Itoh K, Ueno Y, Shinki T, Gillespie MT, Martin TJ, Higashio K, Suda T. Osteoprotegerin produced by osteoblasts is an important regulator in osteoclast development and function. *Endocrinology*. 2000;141(9):3478-84.
54. Boyce BF, Xing L. Functions of RANKL/RANK/OPG in bone modeling and remodeling. *Archives of biochemistry and biophysics*. 2008;473(2):139-146.
55. Xiong J, O'Brien CA. Osteocyte RANKL: New Insights into the Control of Bone Remodeling. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2012;27(3):499-505.
56. Nakashima T, Hayashi M, Fukunaga T, et al. Evidence for osteocyte regulation of bone homeostasis through RANKL expression. *Nat Med* 2011;17:1231–4.
57. Graham JM, Ayati BP, Holstein SA, Martin JA. The role of osteocytes in targeted bone remodeling: a mathematical model. *PLoS One* 2013;8:e63884
58. Cao Z et al. Genetic Evidence for the Vital Function of Osterix in Cementogenesis. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2012;27(5):1080-1092.
59. Nociti FH, Jr., Berry JE, Foster BL, Gurley KA, Kingsley DM, Takata T, et al. Cementum: a phosphate-sensitive tissue. *J Dent Res*. 2002;81:817-21.
60. Foster BL, Nagatomo KJ, Bamashmous SO, Tompkins KA, Fong H, Dunn D, et al. The progressive ankylosis protein regulates cementum apposition and extracellular matrix composition. *Cells Tissues Organs*. 2011;194:382-405.

## FIGURES




**Figure 1.** Schematic illustration of parameters assessed: cellular cementum area (white dotted line area), cementocyte number (white points), cellular cementum density (*e.g.*, number of cementocyte /cellular cementum area - purple area) and cementocyte body size (white points area). Yellow lines indicate as root dentin was measured (166  $\mu\text{m}$  length) to guide the measurement on cellular cementum at the final portion of mandibular 1<sup>st</sup> molar.



**Figure 2.** Increased cementum under experimentally-induced dental cementum apposition model. (A) During early cementum formation (6 days) mandibular first molars featured similar volume in both groups, whereas at 21 days was observed increased area between the groups, however experimental group presented higher values. (B, C) Confocal microscope images, labeling with Fluorescein Isothiocyanate x5 (FITC). In the apex region of first mandibular molars was observed increased dental cementum area in EIA group.


Abbreviations: CE- Dental Cementum; PDL - periodontal ligament; DE- dentin; AB- alveolar bone.

\*Uppercase letter indicate comparisons over time in the same treatment group (intragroup). Lowercase letter indicate treatment comparison at each time point (intergroup). Same letters indicate no significant differences, while values with different letters were significantly different ( $p < 0.05$ ) as tested by ANOVA followed by the Tukey test. Graphs show mean SD values for samples ( $n=5$ ).



**Figure 3.** Experimentally-induced dental cementum apposition model does not affect the cementocyte number. (A) A quantitative approach was performed and cementocytes in a previously established area were counted. Increased cell counting at 21 days was observed, however experimental group value was not significantly different compared to control. (B) Relation of cementocyte number in pre-established area featured decreased significant difference under EIA conditions at 21 days. Suggesting that the amount of matrix formation is responsible for this inversely proportional relationship.


\*Uppercase letter indicate comparisons over time in the same treatment group (intragroup). Lowercase letter indicate treatment comparison at each time point (intergroup). Same letters indicate no significant differences, while values with different letters were significantly different ( $p < 0.05$ ) as tested by ANOVA followed by the Tukey test. Graphs show mean SD values for samples ( $n=5$ ).



**Figure 4.** Increased cell area under experimentally-induced dental cementum apposition model. (A) Experimentally-induced dental cementum apposition model does not affect cell area at 6 days. On the other hand, 21 days experimental group featured significantly different value compared to control. (B, C) Transmission Electron Microscopy (TEM) images reinforces the findings after measurements in confocal images showing cementocytes increased cellular body area at 21 days x 10000.

\*Uppercase letter indicate comparisons over time in the same treatment group (intragroup). Lowercase letter indicate treatment comparison at each time point (intergroup). Same letters indicate no significant differences, while values with different letters were significantly different ( $p < 0.05$ ) as tested by ANOVA followed by the Tukey test. Graphs show mean SD values for samples ( $n=5$ ).













**Figure 5.** Confocal microscope images, labeling with Fluorescein Isothiocyanate (FITC). (A, E) Section of mandibular first molar x5. (B, F) In the apex region was observed increased dental cementum area of EIA group x20. (C, G) Region of apical cementum at higher magnification, x63 (zoom 1.98). No difference related to dendritic processes both in number and in length was detected neither at 21 days experimental or control group. (D, H) Surrounding alveolar bone x63 (zoom 1.98). Photomicrographs revealed irregularly shaped cementocyte lacunae surrounded by few canaliculi compared to alveolar bone. Therefore, lacuno-canalicular network of the alveolar bone (AB) is better developed than in dental cementum (DC).

Arrows indicate cementocytes lacunae.

Head arrows indicate cellular projections

Abbreviations: CE - Dental Cementum; Ccy - cementocyte, PDL - periodontal ligament; DE - dentin; AB - alveolar bone; Ocy - Osteocyte.


**21 days EIA****21 days Control**

**Figure 6.** Scanning Electron Microscope show dental cementum morphology after experimentally-induced apposition (EIA) at 21 days. (A, B) Transverse section of mice mandibular molar x100. (C, D) Apical cementum at higher magnification x500. SEM images confirm that cementocytes occupying lacunae irregularly shaped and unevenly distributed in the matrix and numerous empty cementocyte lacunae. The two patterns observed were lacunae with cell remnants or empty lacunae. (E, F) Internal lacunae x3500. Cementum near the root surface presented lacunae larger in size in both groups. (G, H) External lacunae x3500. Representative SEM images show remnants cementocytes at 21 days.

Arrows indicate cementocytes lacunae.

Abbreviations: CE- Dental Cementum; PDL- periodontal ligament; DE- dentin; AB- alveolar bone.





**Figure 7.** Transmission Electron Microscopy (TEM) demonstrated changes on cellular ultrastructure of cementoblasts and cementocytes at the apical root of mandibular 1<sup>st</sup> molars in experimentally-induced apposition groups. (A) Cementoblasts at the cementum surface and cementocyte early trapped x3000. (B, C) At higher magnification it is possible to observe prominent and well-defined cell nuclei and abundant cytoplasm with rough endoplasmic reticulum (RER) and numerous ribosomes x12000. (C, F) The control group presented cementoblast with well-defined nuclei, intact nuclear membrane and well-developed rough endoplasmic reticulum (RER) x10000; x20000. (B, D, E) EIA group exhibited cementoblasts with well-defined nuclei and intact nuclear membrane. RER is more developed than the control group, indicating intense protein synthesis x10000; x20000. (G) Cementocytes trapped in the lacunae in different layers x3000. (H, I) Both the experimental and control groups' cementocytes presented smaller size compared to cementoblasts, defined nuclei with intact nuclear membrane and poorly developed rough endoplasmic reticulum X10000. (H) At 21 days we observed contracted cells within lacunae, while the control seemed to fill the lacunae even though cell size under experimentally-induced apposition were larger. Experimentally-induced apposition group featured increased cell area and nuclei x10000. (J) Cementum matrix presented radially arranged fibers bundles x10000. (K, L) TEM show a sharp difference in the cellular nuclei and cell proportion in the experimental group, where nuclei occupy substantial cell area whereas this ratio is lower in the control. Cementocyte present not only larger nuclei but also nuclei have more peripherally condensed chromatin and euchromatin spread in the nuclei than the controls after experimental induction x20000.

Abbreviations: Cmb- cementoblast; Ccy- cementocyte; Nu- nucleus; RER- rough endoplasmic reticulum.

### **3 CONCLUSÃO**

Os dados mostraram que o modelo experimental foi eficiente em estimular a aposição de cimento, uma maior quantidade de área de tecido na região apical foi observada no grupo estimulado em relação ao controle em período mais tardio. Os achados morfológicos e ultraestruturais sugerem que os cementócitos, juntamente com cementoblastos, podem desempenhar um papel importante no controle da aposição experimental da matriz do cimento dental formado.

## REFERÊNCIAS

1. Bartold PM. Turnover in periodontal connective tissues: dynamic homeostasis of cells, collagen and ground substances. *Oral Dis.* 1995;1:238-53.
2. Bellido T. Osteocyte-driven bone remodeling. *Calcif Tissue Int.* 2014;94:25-34.
3. Bosshardt DD, Selvig KA. Dental cementum: the dynamic tissue covering of the root. *Periodontol 2000.* 1997;13:41-75.
4. Bosshardt DD, Schroeder HE. Cementogenesis reviewed: a comparison between human premolars and rodent molars. *Anat Rec.* 1996;245:267-92.
5. Bosshardt DD, Sculean A. Does periodontal tissue regeneration really work? *Periodontol 2000.* 2009;51:208-19.
6. Chu EY, Fong H, Blethen FA, Tompkins KA, Foster BL, Yeh KD, et al. Ablation of systemic phosphate-regulating gene fibroblast growth factor 23 (Fgf23) compromises the dentoalveolar complex. *Anat Rec.* 2010;293:1214-26.
7. Compton JT, Lee FY. A review of osteocyte function and the emerging importance of sclerostin. *J Bone Joint Surg Am.* 2014;96:1659-68.
8. Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr Rev.* 2013;34:658-90.
9. Feng JQ, Huang H, Lu Y, Ye L, Xie Y, Tsutsui TW, et al. The Dentin matrix protein 1 (Dmp1) is specifically expressed in mineralized, but not soft, tissues during development. *J Dent Res.* 2003;82:776-80.
10. Feng JQ, Ward LM, Liu S, Lu Y, Xie Y, Yuan B, et al. Loss of DMP1 causes rickets and osteomalacia and identifies a role for osteocytes in mineral metabolism. *Nat Genet.* 2006;38:1310-5.



11. Foster BL, Popowics TE, Fong HK, Somerman MJ. Advances in defining regulators of cementum development and periodontal regeneration. *Curr Top Dev Biol.* 2007;78:47-126.
12. Foster BL, Nagatomo KJ, Bamashmous SO, Tompkins KA, Fong H, Dunn D, et al. The progressive ankylosis protein regulates cementum apposition and extracellular matrix composition. *Cells Tissues Organs.* 2011;194:382-405.
13. Foster BL, Nagatomo KJ, Nociti FH, Fong H, Dunn D, Tran AB, et al. Central role of pyrophosphate in acellular cementum formation. *PLoS One.* 2012;7:e38393.
14. Foster BL, Nociti FH, Jr., Somerman MJ. Tooth Root Formation. In: Huang GTJ, Thesleff I, editors. *Stem Cells, Craniofacial Development and Regeneration.* 1 ed: Wiley-Blackwell; 2013. p. 153-77.
15. Foster BL, Somerman MJ. Cementum. In: McCauley LK, Somerman MJ, editors. *Mineralized Tissues in Oral and Craniofacial Science: Biological Principles and Clinical Correlates.* 1st ed. Ames, IA: Wiley-Blackwell; 2012. p. 169-92.
16. Gluhak-Heinrich J, Ye L, Bonewald LF, Feng JQ, MacDougall M, Harris SE, et al. Mechanical loading stimulates dentin matrix protein 1 (DMP1) expression in osteocytes in vivo. *J Bone Miner Res.* 2003;18:807-17.
17. Grzesik WJ, Narayanan AS. Cementum and periodontal wound healing and regeneration. *Crit Rev Oral Biol Med.* 2002;13:474-84.
18. Hefti AF. Aspects of cell biology of the normal periodontium. *Periodontol* 2000. 1993;3:64-75.
19. Honasoge M, Rao AD, Rao SD. Sclerostin: recent advances and clinical implications. *Curr Opin Endocrinol Diabetes Obes.* 2014;21:437-46.
20. Ito N, Wijenayaka AR, Prideaux M, Kogawa M, Ormsby RT, Evdokiou A, et al. Regulation of FGF23 expression in IDG-SW3 osteocytes and human bone by pro-inflammatory stimuli. *Mol Cell Endocrinol.* 2015;399:208-18.

21. Jager A, Gotz W, Lossdorfer S, Rath-Deschner B. Localization of SOST/sclerostin in cementocytes in vivo and in mineralizing periodontal ligament cells in vitro. *J Periodontal Res.* 2010;45:246-54.
22. Kondoh S, Inoue K, Igarashi K, Sugizaki H, Shirode-Fukuda Y, Inoue E, et al. Estrogen receptor alpha in osteocytes regulates trabecular bone formation in female mice. *Bone.* 2014; 60:68-77.
23. Kuchler U, Schwarze UY, Dobsak T, Heimel P, Bosshardt DD, Kneissel M, et al. Dental and periodontal phenotype in sclerostin knockout mice. *Int J Oral Sci.* 2014;6:70-6.
24. Lau KH, Baylink DJ, Zhou XD, Rodriguez D, Bonewald LF, Li Z, et al. Osteocyte-derived insulin-like growth factor I is essential for determining bone mechanosensitivity. *Am J Physiol Endocrinol Metab.* 2013;305:E271-81.
25. Lee JW, Yamaguchi A, Iimura T. Functional heterogeneity of osteocytes in FGF23 production: the possible involvement of DMP1 as a direct negative regulator. *Bonekey Rep.* 2014;3:543.
26. Lehnen SD, Gotz W, Baxmann M, Jager A. Immunohistochemical evidence for sclerostin during cementogenesis in mice. *Ann Anat.* 2012;194:415-21.
27. Lindberg I, Pang HW, Stains JP, Clark D, Yang AJ, Bonewald L, et al. FGF23 is endogenously phosphorylated in bone cells. *J Bone Miner Res.* 2014.
28. Luan X, Ito Y, Holliday S, Walker C, Daniel J, Galang TM, et al. Extracellular matrix-mediated tissue remodeling following axial movement of teeth. *J Histochem Cytochem.* 2007;55:127-40.
29. Nociti FH, Jr., Foster BL, Tran AB, Dunn D, Presland RB, Wang L, et al. Vitamin D represses dentin matrix protein 1 in cementoblasts and osteocytes. *J Dent Res.* 2014;93:148-54.
30. Nociti FH, Jr., Berry JE, Foster BL, Gurley KA, Kingsley DM, Takata T, et al. Cementum: a phosphate-sensitive tissue. *J Dent Res.* 2002;81:817-21.

31. Sapir-Koren R, Livshits G. Osteocyte control of bone remodeling: is sclerostin a key molecular coordinator of the balanced bone resorption-formation cycles? *Osteoporos Int.* 2014;25:2685-700.
32. Sapir-Koren R, Livshits G. Bone mineralization is regulated by signaling cross talk between molecular factors of local and systemic origin: The role of fibroblast growth factor 23. *Biofactors.* 2014.
33. Salmon CR, Tomazela DM, Ruiz KG, Foster BL, Paes Leme AF, Sallum EA, et al. Proteomic analysis of human dental cementum and alveolar bone. *J Proteomics.* 2013;91C:544-55.
34. Saygin NE, Giannobile WV, Somerman MJ. Molecular and cell biology of cementum. *Periodontol 2000.* 2000;24:73-98.
35. St John HC, Meyer MB, Benkusky NA, Carlson AH, Prideaux M, Bonewald LF, et al. The parathyroid hormone-regulated transcriptome in osteocytes: Parallel actions with 1,25-dihydroxyvitamin D to oppose gene expression changes during differentiation and to promote mature cell function. *Bone.* 2014;72C:81-91.
36. Toyosawa S, Oya K, Sato S, Ishida K. Osteocyte and DMP1. *Clin Calcium.* 2012;22:713-20.
37. Zhao S, Zhang YK, Harris S, Ahuja SS, Bonewald LF. MLO-Y4 osteocyte-like cells support osteoclast formation and activation. *J Bone Miner Res.* 2002;17:2068-79.

**ANEXO 1.** Certificado de aprovação do Comitê de Ética no Uso de Animais – Universidade Estadual de Campinas/UNICAMP.



CEUA/Unicamp

**CERTIFICADO**

Certificamos que o projeto intitulado "**Determinação do papel dos cementócitos na hemostasia do cemento dental**", protocolo nº **3788-1**, sob a responsabilidade de **Prof. Dr. Francisco Humberto Nociti Júnior / Elis Janaina Lira dos Santos**, que envolve a produção, manutenção e/ou utilização de animais pertencentes ao filo *Chordata*, subfilo *Vertebrata* (exceto o homem) para fins de pesquisa científica ou ensino, encontra-se de acordo com os preceitos da **LEI Nº 11.794, DE 8 DE OUTUBRO DE 2008**, que estabelece procedimentos para o uso científico de animais e do **DECRETO Nº 6.899, DE 15 DE JULHO DE 2009**, e com as normas editadas pelo **Conselho Nacional de Controle da Experimentação Animal - CONCEA**, e foi aprovado pela **Comissão de Ética no Uso de Animais da Universidade Estadual de Campinas - CEUA/UNICAMP**, em **14 de maio de 2015**.

**Vigência do projeto:** **07/2015-02/2015**

**Espécie/Linhagem:** **Camundongos Heterogenéticos / Unib:SW (Swiss)**

**No. de animais:** **50**

**Peso/Idade:** **35 dias/10gr**

**Sexo:** **machos**

**Origem:** **CEMIB/UNICAMP**

A aprovação pela CEUA/UNICAMP não dispensa autorização prévia junto ao **IBAMA**, **SISBIO** ou **CIBio**.

Campinas, **14 de maio de 2015**.

Prof. Dr. Alexandre Leite Rodrigues de Oliveira  
Presidente

Fátima Alonso  
Secretária Executiva

CEUA/UNICAMP  
Caixa Postal 6109  
13083-970 Campinas, SP – Brasil

Telefone: (19) 3521-6359  
E-mail: [comisib@unicamp.br](mailto:comisib@unicamp.br)  
<http://www.ib.unicamp.br/ceea/>

**ANEXO 2.** Comprovante de submissão do artigo.

● **Brazilian Journal of Oral Sciences** <onbehalfof+brjorals+fop.unicamp.br@manuscriptcentral.com> Fev 13 em 2:35 PM  
Para elis.lira@yahoo.com.br  
CC elis.lira@yahoo.com.br, crisr.salmon@gmail.com, mandinha\_bandeira@hotmail.com, enilson@unicamp.br,  
casati@unicamp.br, e 4 mais...

13-Feb-2017

Dear Miss Santos:

Your manuscript entitled "ULTRASTRUCTURAL ANALYSIS OF CEMENTOCYTES UNDER EXPERIMENTALLY-INDUCED DENTAL CEMENTUM APPPOSITION" has been successfully submitted online and is presently being given full consideration for publication in the Brazilian Journal of Oral Sciences.

Your manuscript ID is BJOS-2017-0020.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at <https://mc04.manuscriptcentral.com/bjos-scielo> and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to <https://mc04.manuscriptcentral.com/bjos-scielo>.

Thank you for submitting your manuscript to the Brazilian Journal of Oral Sciences.

Sincerely,  
Brazilian Journal of Oral Sciences Editorial Office